

# Finitistic and Representation Dimensions

Jiaqun WEI \*

Department of Mathematics, Nanjing Normal University

Nanjing 210097, P.R.China

Email: weijiaqun@njnu.edu.cn

## Abstract

Let  $A$  be an artin algebra with representation dimension not more than 3. Assume that  ${}_A V$  is an Auslander generator and  $M \in \text{add}_A V$  (for example,  $M \in \text{add}_A(A \oplus \mathbf{D}A)$ ), we indicate that both  $\text{findim}(\text{End}_A M)$  and  $\text{findim}(\text{End}_A M)^{op}$  are finite and consequently, the Gorenstein Symmetry Conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for  $\text{End}_A M$ . In particular, we see that the finitistic dimension conjecture and all the above conjectures hold for artin algebras which can be realized as endomorphism algebras of modules over representation-finite algebras. It is also shown that if every quasi-hereditary algebra has a left idealized extension which is a monomial algebra or an algebra whose representation dimension is not more than 3, then the finitistic dimension conjecture holds.

*MSC:* 16E05; 16E10; 16G20

*Key words:* finitistic dimension; representation dimension; representation-finite algebra

## 1 Introduction

Throughout the paper, we work on artin algebras and finitely generated left modules.

Let  $A$  be an artin algebra. Recall that the little finitistic dimension of  $A$ , denoted by  $\text{findim} A$ , is defined to be the supremum of the projective dimensions of all finitely generated modules of finite projective dimension. Similarly, the big finitistic dimension  $\text{Findim} A$  is defined, allowing arbitrary  $A$ -modules.

In 1960, Bass [4] formulated two so-called finitistic dimension conjectures. The first one asserts that  $\text{findim} A = \text{Findim} A$ , while the second one claims that  $\text{findim} A < \infty$ .

---

\*Supported by the National Science Foundation of China (No.10601024)

It is known that the first finitistic dimension conjecture fails in general [19] and the differences can even be arbitrarily big [14]. However, the second finitistic dimension conjecture is still open. This conjecture is also related to many other homological conjectures (e.g., the Gorenstein Symmetry Conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture) and attracts many algebraists, see for instance [1, 8, 16, 20 etc.].

Only few classes of algebras was known to have finite finitistic dimension, see for instance [6, 7, 8, etc.]. In 1991, Auslander and Reiten [3] proved that  $\text{findim} A < \infty$  if  $\mathcal{P}^{<\infty}$ , the category of all  $A$ -modules of finite projective dimension, is contravariantly finite. However, Igusa-Smalø-Todorov [9] provided an example of artin algebras over which  $\mathcal{P}^{<\infty}$  is not contravariantly finite.

In 2002, Angeleri-Hügel and Trlifaj [1] obtained a sufficient and necessary condition for the finiteness of  $\text{findim} A$ , using the theory of infinitely generated tilting modules. They proved that  $\text{findim} A$  is finite if and only if there exists a special tilting  $A$ -module (maybe infinitely generated).

In the same year, Igusa and Todorov [10] presented a good way to prove the (second) finitistic dimension conjecture. In particular, they proved that  $\text{findim} A$  is finite provided that the representation dimension  $\text{repdim} A$ , another important homological invariant introduced by Auslander [2], is not more than 3. Recall that  $\text{repdim} A = \inf\{\text{gd}(\text{End}_A V) \mid V \text{ is a generator-cogenerator}\}$ , where  $\text{gd}$  denotes the global dimension and  $\text{End}_A V$  denotes the endomorphism algebra of  ${}_A V$ . It should be noted that, though the representation dimension of an artin algebra is always finite [11], there is no upper bound for the representation dimension of an artin algebra [13].

Using the Igusa-Todorov functor defined in [10], Xi [15, 16, 17] developed some new ideas to prove the finiteness of finitistic dimension of some artin algebras. For example, it was shown in [16] that, if  $A$  is a subalgebra of  $R$  such that  $\text{rad} A$  is a left ideal in  $R$  (in the case  $R$  is called a left idealized extension of  $A$ ), then  $\text{findim} A$  is finite provided that  $\text{repdim} R$  is not more than 3. In addition, it was proved that the finitistic dimension conjecture holds for all artin algebras if and only if the finiteness of  $\text{findim} R$  implies the finiteness of  $\text{findim} A$ , for any pair  $A, R$  such that  $R$  is a left idealized extension of  $A$  [15].

In this note, we will continue the study of the finitistic dimension conjecture following the ideas comes from [10, 15, 16, 17]. Recall that an Auslander generator over the artin algebra  $A$  is a generator-cogenerator  ${}_A V$  such that  $\text{repdim} A = \text{gd}(\text{End}_A V)$ . Note that it is not known if  $\text{findim} A^{op}$  is finite provided that  $\text{findim} A$  is finite in general, where  $A^{op}$  denotes the opposite algebra of  $A$ .

Our main results state as follows.

**Theorem 1.1** *Let  $A$  be an artin algebra with  $\text{repdim} A \leq 3$ . Assume that  ${}_A V$  is an Aus-*

lander generator. Then both  $\text{findim}(\text{End}_A M)$  and  $\text{findim}(\text{End}_A M)^{op}$  are finite, whenever  $M \in \text{add}_A V$ .

Let  $E$  be an artin algebra. Recall the following well known conjectures (see for instance [8, 12], etc.).

**Gorenstein Symmetry Conjecture**  $\text{id}_E E < \infty$  if and only if  $\text{id}(E_E) < \infty$ , where  $\text{id}$  denotes the injective dimension.

**Wakamatsu-tilting Conjecture** Let  ${}_E \omega$  be a Wakamatsu-tilting module. (1) If  $\text{pd}_E \omega < \infty$ , then  $\omega$  is tilting. (2) If  $\text{id}_E \omega < \infty$ , then  $\omega$  is cotilting.

**Generalized Nakayama Conjecture** Each indecomposable injective  $E$ -module occurs as a direct summand in the minimal injective resolution of  ${}_E E$ .

Note that the Gorenstein symmetry conjecture and the Generalized Nakayama conjecture are special cases of the second Wakamatsu-tilting conjecture. Moreover, if the finitistic dimension conjecture for  $E$  and  $E^{op}$ , then all the above conjectures hold.

**Corollary 1.2** *Let  $A$  be an artin algebra with  $\text{repdim} A \leq 3$ . Assume that  ${}_A V$  is an Auslander generator. Then the Gorenstein Symmetry Conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for  $\text{End}_A M$ .*

As a special case of and Theorem 1.1 and Corollary 1.2, we obtain the following result.

**Corollary 1.3** *If  $A = \text{End}_\Lambda M$  for some module  $M$  over a representation-finite algebra  $\Lambda$ , then both  $\text{findim} A$  and  $\text{findim} A^{op}$  are finite. In particular, the Gorenstein Symmetry Conjecture, the Wakamatsu-tilting conjecture and the generalized Nakayama conjecture hold for  $A$ .*

We do not know whether or not every artin algebra can be realized as an endomorphism algebra of some module over a representation-finite algebra. If it is the case, then the above result indicates that the finitistic dimension conjecture holds for all artin algebras.

However, it is well known that every artin algebra can be realized as an endomorphism algebra of a projective and injective module over a quasi-hereditary algebra. Thus the following result shows that, in particular, if every quasi-hereditary algebra has a left idealized extension which is a monomial algebra or an algebra whose representation dimension is not more than 3, then the finitistic dimension conjecture holds.

**Proposition 1.4** *Let  $R$  be a left idealized extension of  $A$ . If  $\text{repdim} R \leq 3$  or  $\Omega_R^2(R\text{-mod})$  is of finite type (for example,  $\text{gd} R \leq 2$ ), then  $\text{findim}(\text{End}_A M)$  is finite, for any projective  $A$ -module  $M$ .*

## 2 Finitistic dimension of endomorphism algebras

Let  $A$  be an artin algebra. We denote by  $A\text{-mod}$  the category of all  $A$ -modules. If  $M$  is an  $A$ -module, we use  $\text{pd}_A M$  to denote the projective dimension of  $M$  and use  $\Omega_A^i M$  to denote the  $i$ -th syzygy of  $M$ . Throughout the note,  $\mathbf{D}$  denotes the usual duality functor between  $A\text{-mod}$  and  $A^{op}\text{-mod}$ .

The following lemma was well-known, see for instance [17].

**Lemma 2.1** *Let  $A$  be an artin algebra and let  $V$  be a generator-cogenerator in  $A\text{-mod}$ . The following are equivalent for a nonnegative integer  $n$ .*

- (1)  $\text{gd}(\text{End}_A V) \leq n + 2$ .
- (2) *For any  $X \in A\text{-mod}$ , there is an exact sequence  $0 \rightarrow V_n \rightarrow \cdots \rightarrow V_1 \rightarrow V_0 \rightarrow X \rightarrow 0$  with each  $V_i \in \text{add}_A V$ , such that the corresponding sequence induced by the functor  $\text{Hom}_A(V, -)$  is also exact.*

The following lemma collects some important properties of the Igusa-Todorov functor introduced in [10].

**Lemma 2.2** *For any artin algebra  $A$ , there is a functor  $\Psi$  which is defined on the objects of  $A\text{-mod}$  and takes nonnegative integers as values, such that*

- (1)  $\Psi(M) = \text{pd}_A M$  provided that  $\text{pd}_A M < \infty$ .
- (2)  $\Psi(X) \leq \Psi(Y)$  whenever  $\text{add}_A X \subseteq \text{add}_A Y$ . The equation holds in case  $\text{add}_A X = \text{add}_A Y$ .
- (3) *If  $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$  is an exact sequence in  $A\text{-mod}$  with  $\text{pd}_A Z < \infty$ , then  $\text{pd}_A Z \leq \Psi(X \oplus Y) + 1$ .*

Let  $A$  be an artin algebra and  $M \in A\text{-mod}$  with  $E = \text{End}_A M$ . Then  $M$  is also a right  $E$ -module. It is well known that  $(M \otimes_E -, \text{Hom}_A(M, -))$  is a pair of adjoint functors and that, for any  $E$ -module  $Y$ , there is a canonical homomorphism  $\sigma_Y : Y \rightarrow \text{Hom}_A(M, M \otimes_E Y)$  defined by  $n \rightarrow [t \rightarrow t \otimes n]$ . It is easy to see that  $\sigma_Y$  is an isomorphism provided that  $Y$  is a projective  $E$ -module.

The following lemma is elementary but essential to prove our results.

**Lemma 2.3** *Let  $M \in A\text{-mod}$  and  $E = \text{End}_A M$ . Then, for any  $X \in E\text{-mod}$ ,  $\Omega_E^2 X \simeq \text{Hom}_A(M, Y)$  for some  $Y \in A\text{-mod}$ .*

**Proof.** Obviously, we have an exact sequence

$$0 \rightarrow \Omega_E^2 X \rightarrow E_1 \rightarrow E_0 \rightarrow X \rightarrow 0$$

with  $E_0, E_1 \in E\text{-mod}$  projective. Applying the functor  $M \otimes_E -$ , we obtain an induced exact sequence

$$0 \rightarrow Y \rightarrow M \otimes_E E_1 \rightarrow M \otimes_E E_0 \rightarrow M \otimes_E X \rightarrow 0,$$

for some  $Y \in A\text{-mod}$ . Now applying the functor  $\text{Hom}_A(M, -)$ , we further have an induced exact sequence

$$0 \rightarrow \text{Hom}_A(M, Y) \rightarrow \text{Hom}_A(M, M \otimes_E E_1) \rightarrow \text{Hom}_A(M, M \otimes_E E_0).$$

Moreover, there is the following commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow & \Omega_E^2 X & \rightarrow & E_1 & \rightarrow & E_0 & \\ & \downarrow \phi & & \downarrow \sigma_{E_1} & & \downarrow \sigma_{E_0} & \\ 0 \rightarrow & \text{Hom}_A(M, Y) & \rightarrow & \text{Hom}_A(M, M \otimes_E E_1) & \rightarrow & \text{Hom}_A(M, M \otimes_E E_0). & \end{array}$$

Since  $E = \text{End}_A M$  and  $E_0, E_1 \in \text{add}_E E$ , the canonical homomorphisms  $\sigma_{E_0}$  and  $\sigma_{E_1}$  are isomorphisms. It follows that  $\Omega_E^2 X \simeq \text{Hom}_A(M, Y)$ .

#

**Theorem 2.4** *Let  $A$  be an artin algebra and  ${}_A V$  be a generator-cogenerator in  $A\text{-mod}$  such that  $\text{gd}(\text{End}_A V) \leq 3$ . Then  $\text{findim} E$  is finite, where  $E = \text{End}_A M$  for some  $M \in \text{add}_A V$ .*

**Proof.** Suppose that  $X \in E\text{-mod}$  and that  $\text{pd}_E X < \infty$ . Then  $\text{pd}_E(\Omega_E^2 X) < \infty$ . Moreover,  $\Omega_E^2 X \simeq \text{Hom}_A(M, Y)$  for some  $Y \in A\text{-mod}$ , by Lemma 2.3. Since  ${}_A V$  is a generator and  $\text{gd}(\text{End}_A V) \leq 3$ , by Lemma 2.1 we obtain an exact sequence

$$0 \rightarrow V_1 \rightarrow V_0 \rightarrow Y \rightarrow 0 \quad (\dagger)$$

with  $V_0, V_1 \in \text{add}_A V$  such that the corresponding sequence induced by the functor  $\text{Hom}_A(V, -)$  is also exact. Note that  $M \in \text{add}_A V$ , so the sequence  $(\dagger)$  also stays exact under the functor  $\text{Hom}_A(M, -)$ . Thus, we have the following exact sequence in  $E\text{-mod}$ :

$$0 \rightarrow \text{Hom}_A(M, V_1) \rightarrow \text{Hom}_A(M, V_0) \rightarrow \text{Hom}_A(M, Y) \rightarrow 0.$$

Now by Lemma 2.2, we deduce that

$$\begin{aligned} \text{pd}_E X &\leq \text{pd}_E(\Omega_E^2 X) + 2 \\ &= \text{pd}_E(\text{Hom}_A(M, Y)) + 2 \\ &\leq \Psi(\text{Hom}_A(M, V_0) \oplus \text{Hom}_A(M, V_1)) + 1 + 2 \\ &\leq \Psi(\text{Hom}_A(M, V)) + 1 + 2 < \infty. \end{aligned}$$

#

It is not known if  $\text{findim} A^{op}$  is finite provided that  $\text{findim} A$  is finite in general. For an artin algebra  $A$  with  $\text{repdim} A \leq 3$ , it is known that  $\text{findim} A$  is finite [11] and since  $\text{repdim} A = \text{repdim} A^{op}$ ,  $\text{findim} A^{op}$  is also finite in the case. The following corollary is also a generalization of this fact.

**Corollary 2.5** *Let  $A$  be an artin algebra with  $\text{repdim} A \leq 3$ . Assume that  ${}_A V$  is an Auslander generator. If  $M \in \text{add}_A V$ , then both  $\text{findim}(\text{End}_A M)$  and  $\text{findim}(\text{End}_A M)^{op}$  are finite. In particular, the result holds for any  ${}_A M \in \text{add}_A(A \oplus \mathbf{D}A)$ .*

**Proof.** If  ${}_A V$  is an Auslander generator in  $A\text{-mod}$ , then  ${}_{A^{op}} \mathbf{D}V$  is an Auslander generator in  $A^{op}\text{-mod}$ . Moreover, if  $M \in \text{add}_A V$ , then  $\mathbf{D}M \in \text{add}_{A^{op}} \mathbf{D}V$ . Obviously,  $(\text{End}_A M)^{op} \simeq \text{End}_{A^{op}} \mathbf{D}M$ . Now the conclusion follows from Theorem 2.4.

#

Since all representation-finite algebras have representation dimension 2, we obtain the following corollary.

**Corollary 2.6** *If  $A = \text{End}_\Lambda M$  for some representation-finite algebra  $\Lambda$  and some  $M \in \Lambda\text{-mod}$ , then both  $\text{findim} A$  and  $\text{findim} A^{op}$  are finite.*

**Proof.** Since  $\Lambda$  is of finite representation type, the module  $V = \bigoplus_{i=1}^n V_i$  is an Auslander generator, where  $\{V_i, i = 1, \dots, n\}$  is the representation set of all indecomposable  $\Lambda$ -modules. Now the conclusion follows from Corollary 2.5.

#

The above result suggests a strong connection between the finitistic dimension conjecture and the following endomorphism algebra realization problem of artin algebras.

**Problem 1** Can all artin algebras be realized as endomorphism algebras of modules over representation-finite algebras? If not, what algebra has such a realization?

Obviously, the affirmative answer to the first question implies the finitistic dimension conjecture holds, by Corollary 2.6.

In contrast, it is well known that every artin algebra can be realized as an endomorphism algebra of a projective and injective module over a quasi-hereditary algebra.

Let  $A, R$  be both Artin algebras. Following [15], we say that  $R$  is a left idealized extension of  $A$  if  $A \subseteq R$  has the same identity and  $\text{rad} A$  is a left ideal in  $R$ .

**Proposition 2.7** *Let  $R$  be a left idealized extension of  $A$ . If  $\text{repdim} R \leq 3$  or  $\Omega_R^2(R\text{-mod})$  is of finite type (for example,  $\text{gd} R \leq 2$ ), then  $\text{findim} E$  is finite, where  $E = \text{End}_A M$  for some projective  $A$ -module  $M$ .*

**Proof.** Suppose that  $X \in E\text{-mod}$  and that  $\text{pd}_E X < \infty$ . Then  $\text{pd}_E(\Omega_E^2 X) < \infty$ . Moreover,  $\Omega_E^2 X \simeq \text{Hom}_A(M, Y)$  for some  $Y \in A\text{-mod}$ , by Lemma 2.3. Since  ${}_A M$  is projective, the proof of Lemma 2.3 in fact indicates that  $Y \in \Omega_A^2(A\text{-mod})$ . Then  $Y \simeq \Omega_R Z \oplus Q$  as  $R$ -modules for some  $Z, Q \in R\text{-mod}$  with  $Q$  projective, by [15, Erratum]. If  $\text{repdim} R \leq 3$  or  $\Omega_R^2(R\text{-mod})$  is of finite type, we can obtain an exact sequence of  $R$ -modules

$$0 \rightarrow V_1 \rightarrow V_0 \rightarrow Y \rightarrow 0 \quad (\ddagger)$$

with  $V_0, V_1 \in \text{add}_R V$ . Here  ${}_R V$  is an Auslander generator in case  $\text{repdim} R \leq 3$  or  $V = R \oplus N$  in case  $\Omega_R^2(R\text{-mod})$  is of finite type, where  $N$  is the direct sum of nonisomorphic indecomposable  $R$ -modules appeared in  $\Omega_R^2(R\text{-mod})$ . The exact sequence  $(\ddagger)$  then restricts to an exact sequence in  $A\text{-mod}$ . Since  ${}_A M$  is projective, we have the following exact sequence in  $E\text{-mod}$ :

$$0 \rightarrow \text{Hom}_A(M, V_1) \rightarrow \text{Hom}_A(M, V_0) \rightarrow \text{Hom}_A(M, Y) \rightarrow 0.$$

Now by Lemma 2.2, we deduce that

$$\begin{aligned} \text{pd}_E X &\leq \text{pd}_E(\Omega_E^2 X) + 2 \\ &= \text{pd}_E(\text{Hom}_A(M, Y)) + 2 \\ &\leq \Psi(\text{Hom}_A(M, V_0) \oplus \text{Hom}_A(M, V_1)) + 1 + 2 \\ &\leq \Psi(\text{Hom}_A(M, V)) + 1 + 2 < \infty. \end{aligned}$$

#

The above result has the following corollary which contains [18, Corollary 2.15], where the result was proved under an additional condition.

**Corollary 2.8** *If  $A$  has a left idealized extension which is a monomial algebra, then  $\text{findim} E$  is finite, where  $E = \text{End}_A M$  for some projective  $A$ -module  $M$ .*

**Proof.** It was noted that  $\Omega_R^2(R\text{-mod})$  is of finite type in case  $R$  is a finite dimensional monomial relation algebra, see [5, Example 2.3(a)] (also [19, Theorem A]). Thus the conclusion follows from Proposition 2.7.

#

**Problem 2** Does every quasi-hereditary algebra have a left idealized extension which is a monomial algebra or an algebra whose representation dimension is not more than 3?

Of course, the affirmative answer to the problem also implies the finitistic dimension conjecture holds.

## References

- [1] Angeleri-Hügel L. and Trlifaj J., Tilting theory and the finitistic dimension conjecture, Trans. Amer. Math. Soc. 354 (11) (2002), 4345-4358.

- [2] Auslander M., Representation dimension of artin algebras, Queen Mary College Mathematics Notes, London (1971) (Republished in: M. Auslander, Selected works of Maurice Auslander. Part 1. Edited and with a foreword by Idun Reiten, Sverre O. Smalø and Øyvind Solberg. American Mathematical Society, Providence, RI, 1999. xxii+895 pp.)
- [3] Auslander M. and Reiten I., Applications of contravariantly finite subcategories, *Adv. Math.* 86 (1991), 111-152.
- [4] Bass H., Finitistic dimension and a homological generalization of semiprimary rings, *Trans AMS* 95 (1960), 466-488.
- [5] Goodearl K. and Zimmermann-huisgen B., Repetitive resolutions over classical orders and finite dimensional algebras, in *Algebras and Modules II* (Reiten I., Smalø S.O., Solberg O., Eds.) *Canad. Math. Soc. Conf. Proc. Series* 24 (1998) 205-225.
- [6] Green E. L., Kirmann E. and Kuzmanovich J., Finitistic dimensions of finite monomial algebras, *J. Algebra* 136 (1991), 37-50.
- [7] Green E. L. and Zimmermann-huisgen B., Finitistic dimension of artinian rings with vanishing radical cube, *Math. Z.* 206 (1991), 505-526.
- [8] Happel D., Homological conjectures in representation theory of finite-dimensional algebras, *Sherbrook Lecture Notes Series* (1991), available from <http://www.math.ntnu.no/oyvinso/Nordfjordeid/Program/references.html>.
- [9] [IST] Igusa K., Smalø S. and Todorov, T., Finite projectivity and contravariant finiteness, *Proc. AMS.* 109 (1990), 937-941.
- [10] Igusa K. and Todorov G., On the finitistic global dimension conjecture for artin algebras. In: *Representations of algebras and related topics*, 201-204. *Fields Inst. Commun.* 45, Amer. Math. Soc., Providence, RI, 2005.
- [11] Iyama O., Finiteness of representation dimension, *Proc. AMS.* 131 (2003), 1011C1014.
- [12] Mantese F. and Reiten I., Wakamatsu Tilting modules, *J. Algebra* 278 (2004), 532-552.
- [13] Rouquier R., Dimensions of triangulated categories. *arXiv:math.CT/0310134 v3*.
- [14] Smalø S. O., Homological differences between finite and infinite dimensional representations of algebras, *Trends in Math.*, pp. 81-93, Birkhäuser, Basel, 2000.
- [15] Xi C., On the finitistic dimension conjecture I: related to representation-finite algebras, *J. Pure and Appl. Algebra* 193 (2004), 287-305. Erratum: *J. Pure and Appl. Algebra* 202 (1-3) (2005), 325-328.
- [16] Xi C., On the finitistic dimension conjecture II: related to finite global dimension, *Adv. Math.* 201 (2006), 116-142.
- [17] Xi C., On the finitistic dimension conjecture III: related to the pair  $eAe \subseteq A$ , *J. Algebra* 319 (2008), 3666C3688.
- [18] Xi C. and Xu D., On the finitistic dimension conjecture IV: related to relatively projective modules, manuscript.
- [19] Zimmermann-Huisgen B., Homological domino effects and the first finistic demension conjecture, *Invent. Math.* 108 (1992), 369-383.
- [20] Zimmermann-Huisgen B., The finitistic dimension conjectures - A tale of 3.5 decades, In: *Abelian Groups and Modules*, 501-517, Kluwer, Dordrecht, 1995.